

Table X-1
KAPOSI'S SARCOMA

NUMBER OF CASES

By SEER Location and Year of Diagnosis, 1973-94

All Races, Males, All Ages

SEER Location	Year of Diagnosis															
	1973-79	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994
All Areas [⊕]	174	34	44	109	217	367	517	723	853	1015	1110	1098	1131	1067	964	855
All Areas Excluding SF-Oakland SMSA	130	23	21	35	50	103	153	197	233	353	417	424	411	485	469	396
San Francisco-Oakland SMSA	44	11	23	74	167	264	364	526	620	662	693	674	720	582	495	459
San Francisco City/County	17	6	14	61	140	227	313	441	512	546	532	525	570	426	360	329

AGE-ADJUSTED CANCER INCIDENCE RATES[§]

By SEER Location and Year of Diagnosis, 1973-94

All Races, Males, All Ages

SEER Location	Year of Diagnosis															
	1973-79	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994
All Areas [⊕]	0.3	0.4	0.4	0.9	1.7	2.8	3.8	5.1	6.0	6.9	7.5	7.2	7.4	6.8	6.1	5.4
All Areas Excluding SF-Oakland SMSA	0.3	0.3	0.3	0.4	0.5	1.0	1.4	1.7	2.0	3.0	3.4	3.4	3.2	3.7	3.5	2.9
San Francisco-Oakland SMSA	0.5	0.7	1.3	3.5	7.3	11.5	15.7	21.6	25.4	26.1	27.2	25.7	27.8	22.1	18.4	17.4
San Francisco City/County	0.6	1.5	3.3	12.4	26.7	43.2	59.4	79.7	92.6	97.8	97.1	94.2	103.9	77.8	64.5	60.8

§ SEER Program. Rates are per 100,000 males and are age-adjusted to the 1970 U.S. standard population. Each rate has been age-adjusted by 5-year age groups.

⊕ All Areas refers to all SEER data collection areas. The San Francisco-Oakland SMSA represents 15% of all persons in the SEER areas. For this age group, all SEER areas may not be generalizable to the United States for Kaposi's Sarcoma.

Table X-2
KAPOSI'S SARCOMA

NUMBER OF CASES

By SEER Location and Year of Diagnosis, 1973-94

All Races, Males, Ages 20-54

SEER Location	Year of Diagnosis															
	1973-79	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994
All Areas⊕	18	5	20	75	178	328	475	682	799	944	1043	1037	1063	988	908	808
All Areas Excluding SF-Oakland SMSA	14	<4	4	9	25	77	128	174	205	311	378	388	376	452	436	376
San Francisco-Oakland SMSA	4	4	16	66	153	251	347	508	594	633	665	649	687	536	472	432
San Francisco City/County	<4	4	13	59	134	220	300	428	494	527	515	509	548	396	347	311

AGE-ADJUSTED CANCER INCIDENCE RATES§

By SEER Location and Year of Diagnosis, 1973-94

All Races, Males, Ages 20-54

SEER Location	Year of Diagnosis															
	1973-79	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994
All Areas⊕	0.1	0.1	0.4	1.3	3.0	5.6	7.9	11.0	12.7	14.5	15.8	15.4	15.6	14.0	12.9	11.4
All Areas Excluding SF-Oakland SMSA	0.1	-	0.1	0.2	0.5	1.6	2.5	3.3	3.8	5.7	6.8	6.9	6.5	7.7	7.4	6.3
San Francisco-Oakland SMSA	0.1	0.5	1.9	6.9	14.9	24.9	34.0	47.5	55.3	56.3	59.0	56.1	59.7	44.6	39.6	36.4
San Francisco City/County	-	2.4	7.2	27.7	58.3	96.4	129.6	176.5	203.4	214.8	213.2	207.8	226.5	159.3	141.4	128.7

§ SEER Program. Rates are per 100,000 males aged 20-54 and are age-adjusted to the 1970 U.S. standard population. Each rate has been age-adjusted by 5-year age groups.

⊕ All Areas refers to all SEER data collection areas. The San Francisco-Oakland SMSA represents 15% of all persons in the SEER areas. For this age group, all SEER areas may not be generalizable to the United States for Kaposi's Sarcoma.

Table X-3
KAPOSI'S SARCOMA

NUMBER OF CASES

By SEER Location and Year of Diagnosis, 1973-94

All Races, Males, Ages 55+

SEER Location	Year of Diagnosis															
	1973-79	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994
All Areas [⊕]	156	29	24	34	39	38	42	39	54	70	66	61	68	78	56	47
All Areas Excluding SF-Oakland SMSA	116	22	17	26	25	26	25	22	28	42	39	36	35	32	33	20
San Francisco-Oakland SMSA	40	7	7	8	14	12	17	17	26	28	27	25	33	46	23	27
San Francisco City/County	16	<4	<4	<4	6	7	13	13	18	19	16	16	22	30	13	18

AGE-ADJUSTED CANCER INCIDENCE RATES[§]

By SEER Location and Year of Diagnosis, 1973-94


All Races, Males, Ages 55+

SEER Location	Year of Diagnosis															
	1973-79	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994
All Areas [⊕]	1.5	1.7	1.4	1.9	2.1	2.0	2.2	2.0	2.7	3.5	3.4	3.0	3.4	4.0	2.7	2.3
All Areas Excluding SF-Oakland SMSA	1.3	1.6	1.2	1.7	1.6	1.6	1.6	1.3	1.7	2.5	2.4	2.0	2.1	1.9	1.8	1.1
San Francisco-Oakland SMSA	2.3	2.7	2.4	2.7	4.8	4.0	5.4	5.5	8.3	9.0	8.9	8.1	10.8	15.3	7.2	8.9
San Francisco City/County	2.9	-	-	-	8.4	9.0	18.7	19.0	25.7	27.5	24.7	24.4	32.9	48.0	18.8	28.1

§ SEER Program. Rates are per 100,000 males aged 55+ and are age-adjusted to the 1970 U.S. standard population. Each rate has been age-adjusted by 5-year age groups.


⊕ All Areas refers to all SEER data collection areas. The San Francisco-Oakland SMSA represents 15% of all persons in the SEER areas. For this age group, all SEER areas may not be generalizable to the United States for Kaposi's Sarcoma.

All Cancer Sites vs. Kaposi's Sarcoma SEER Incidence Rates, 1973-1994 Males, Ages 20-54




Age-adjusted to 1970 Standard

San Francisco/Oakland Ages 20-54, All Males, 1975-94



All SEER excluding San Francisco/Oakland Ages 20-54, All Males, 1975-94



*Kaposi's excluded from Oral, Colorec., Lung, Melanoma, Testis, and NHL.