

Table I-20
US AND SEER DEATH RATES BY PRIMARY CANCER SITE AND RACE/ETHNICITY, 2000-2004

		Total United States ^a							SEER 17 Areas ^{ab}						
Site		Total	White	Black	AI/AN ^c	API ^d	Hisp ^e	W-NHisp ^e	Total	White	Black	AI/AN ^c	API ^d	Hisp ^e	W-NHisp ^e
All Sites	Both Sexes	192.7	190.7	238.8	160.4	115.5	129.1	194.7	186.1	187.0	240.0	143.7	122.9	127.9	193.3
	Male	238.7	234.7	321.8	187.9	141.7	162.2	239.2	226.7	226.2	316.0	168.6	152.4	157.3	233.2
	Female	162.2	161.4	189.3	141.2	96.7	106.7	165.1	159.2	161.2	193.8	125.8	101.6	108.1	167.0
Oral Cavity & Pharynx	Both Sexes	2.7	2.5	3.9	2.5	2.3	1.7	2.6	2.7	2.6	3.9	3.0	2.5	1.6	2.7
	Male	4.1	3.8	6.8	3.8	3.4	2.8	3.9	4.0	3.8	6.6	3.8	3.7	2.6	3.9
	Female	1.5	1.5	1.7	1.5	1.3	0.8	1.5	1.6	1.6	1.9	2.4	1.4	0.9	1.6
Esophagus	Both Sexes	4.4	4.3	6.0	3.8	1.8	2.4	4.5	4.1	4.2	5.2	4.1	1.9	2.3	4.4
	Male	7.8	7.7	10.2	6.7	3.1	4.2	7.9	7.2	7.4	9.0	7.2	3.3	4.2	7.8
	Female	1.8	1.7	3.0	1.5	0.8	1.0	1.7	1.7	1.6	2.5	-	0.9	0.9	1.7
Stomach	Both Sexes	4.2	3.7	8.2	7.2	8.0	6.8	3.5	4.7	4.1	8.5	7.8	8.3	7.3	3.6
	Male	5.9	5.2	11.9	9.6	10.5	9.1	4.9	6.6	5.7	12.2	11.0	11.1	9.9	5.1
	Female	3.0	2.6	5.8	5.5	6.2	5.1	2.4	3.4	2.9	6.0	5.7	6.2	5.6	2.5
Colon & Rectum	Both Sexes	19.4	18.9	26.7	17.0	12.3	13.6	19.2	18.7	18.4	26.9	16.6	13.2	12.7	18.9
	Male	23.5	22.9	32.7	20.6	15.0	17.0	23.3	22.3	22.0	32.0	19.7	16.5	16.0	22.6
	Female	16.4	15.9	22.9	14.3	10.3	11.1	16.2	15.9	15.6	23.6	14.5	10.8	10.2	16.1
Liver & Intrahepatic Bile Duct	Both Sexes	4.9	4.5	6.5	8.4	10.6	7.6	4.2	5.4	4.8	6.8	8.7	10.8	7.6	4.4
	Male	7.1	6.5	10.0	10.7	15.5	10.8	6.1	7.9	6.8	10.8	11.0	15.7	10.6	6.3
	Female	3.1	2.8	3.9	6.4	6.7	5.0	2.7	3.4	3.1	3.8	6.9	6.8	5.1	2.8
Pancreas	Both Sexes	10.6	10.4	13.8	7.5	7.4	8.3	10.5	10.5	10.5	14.1	7.2	7.8	8.4	10.7
	Male	12.2	12.0	15.5	7.6	7.9	9.1	12.2	11.9	12.0	15.2	7.6	8.5	8.6	12.3
	Female	9.2	9.0	12.4	7.3	6.9	7.5	9.1	9.4	9.2	13.0	6.9	7.3	8.1	9.3
Larynx	Both Sexes	1.3	1.2	2.5	1.1	0.4	0.9	1.2	1.2	1.1	2.6	-	0.4	0.8	1.2
	Male	2.4	2.2	5.0	1.9	0.8	1.9	2.2	2.2	2.0	5.3	-	0.9	1.7	2.1
	Female	0.5	0.5	0.8	-	0.1	0.2	0.5	0.4	0.4	0.8	-	-	0.1	0.5
Lung & Bronchus	Both Sexes	54.7	55.0	62.0	39.9	26.9	23.6	57.4	50.5	51.4	62.8	30.9	28.7	22.2	54.9
	Male	73.4	72.6	95.8	49.6	38.3	36.0	75.4	65.9	65.8	93.7	39.9	41.5	32.8	69.6
	Female	41.1	42.1	39.8	32.7	18.5	14.6	44.3	39.3	41.1	42.3	24.1	19.2	14.7	44.4
Melanoma of the Skin	Both Sexes	2.6	3.0	0.4	1.0	0.3	0.7	3.2	2.5	3.0	0.4	-	0.4	0.7	3.3
	Male	3.9	4.3	0.5	1.3	0.4	0.9	4.7	3.7	4.4	0.5	-	0.5	0.9	4.9
	Female	1.7	2.0	0.4	0.7	0.3	0.6	2.1	1.6	1.9	0.4	-	0.3	0.6	2.1
Breast	Female	25.5	25.0	33.8	16.1	12.6	16.1	25.5	25.3	25.3	34.9	13.6	13.9	15.7	26.4

- ^a NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- ^b The SEER 17 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey.
- ^c Rates for American Indian/Alaska Native (AI/AN) are based on the CHSDA (Contract Health Service Delivery Area) counties.
- ^d Asian/Pacific Islander.
- ^e Hispanic (Hisp) and White Non-Hispanic (W-NHisp) are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Mortality data for Hispanics and White Non-Hispanics do not include cases from Minnesota, New Hampshire, and North Dakota.
- Statistic could not be calculated due to less than 16 cases in the time interval.

Table I-20 - continued
 US AND SEER DEATH RATES BY PRIMARY CANCER SITE AND RACE/ETHNICITY, 2000-2004

		Total United States ^a							SEER 17 Areas ^{ab}						
Site		Total	White	Black	AI/AN ^c	API ^d	Hisp ^e	W-NHisp ^e	Total	White	Black	AI/AN ^c	API ^d	Hisp ^e	W-NHisp ^e
Cervix	Female	2.6	2.3	4.9	4.0	2.4	3.3	2.2	2.5	2.3	4.3	2.6	2.5	3.4	2.1
Corpus & Uterus, NOS	Female	4.1	3.9	7.1	2.9	2.4	3.1	3.9	4.1	4.0	7.0	2.6	2.6	3.0	4.0
Ovary	Female	8.9	9.2	7.4	7.1	4.8	6.1	9.5	9.1	9.6	7.6	7.3	5.2	6.6	10.0
Prostate	Male	27.9	25.6	62.3	21.5	11.3	21.2	25.6	26.9	25.8	57.9	16.4	12.3	21.0	26.1
Testis	Male	0.3	0.3	0.2	-	0.1	0.2	0.3	0.2	0.3	0.1	-	0.1	0.2	0.3
Urinary Bladder	Both Sexes	4.3	4.5	3.7	1.9	1.8	2.4	4.6	4.2	4.4	3.9	1.6	1.9	2.3	4.6
	Male	7.5	7.9	5.3	3.1	2.9	3.9	8.1	7.1	7.6	5.6	-	3.1	3.7	8.0
	Female	2.3	2.3	2.8	1.1	1.0	1.4	2.3	2.2	2.3	2.9	-	1.0	1.4	2.3
Kidney & Renal Pelvis	Both Sexes	4.2	4.3	4.1	6.5	1.7	3.6	4.3	4.0	4.2	4.1	5.0	1.9	3.6	4.2
	Male	6.1	6.2	6.1	9.3	2.4	5.4	6.2	5.9	6.1	6.2	7.6	2.7	5.1	6.2
	Female	2.8	2.8	2.8	4.3	1.1	2.3	2.8	2.6	2.7	2.7	3.0	1.2	2.5	2.7
Brain & Nervous System	Both Sexes	4.4	4.8	2.6	2.1	2.0	2.8	4.9	4.3	4.8	2.7	1.8	2.0	2.9	5.0
	Male	5.4	5.8	3.2	2.6	2.5	3.4	6.0	5.3	5.8	3.2	2.4	2.7	3.5	6.2
	Female	3.6	3.9	2.2	1.8	1.5	2.4	4.0	3.5	3.9	2.2	-	1.5	2.5	4.1
Thyroid	Both Sexes	0.5	0.5	0.5	0.5	0.6	0.6	0.5	0.5	0.5	0.5	-	0.6	0.6	0.5
	Male	0.5	0.5	0.4	-	0.4	0.5	0.5	0.5	0.5	0.4	-	0.4	0.5	0.5
	Female	0.5	0.5	0.5	0.7	0.7	0.6	0.4	0.5	0.5	0.5	-	0.7	0.7	0.5
Hodgkin Lymphoma	Both Sexes	0.5	0.5	0.4	0.3	0.2	0.4	0.5	0.5	0.5	0.4	-	0.2	0.4	0.5
	Male	0.6	0.6	0.5	-	0.3	0.6	0.6	0.6	0.6	0.5	-	0.3	0.5	0.6
	Female	0.4	0.4	0.3	-	0.2	0.3	0.4	0.4	0.4	0.3	-	0.2	0.3	0.4
Non-Hodgkin Lymphoma	Both Sexes	7.6	7.9	5.2	5.5	4.8	5.7	8.0	7.4	7.8	5.3	5.4	5.2	5.7	8.0
	Male	9.6	9.9	6.5	6.1	5.8	6.9	10.1	9.4	9.9	6.8	5.3	6.4	7.0	10.2
	Female	6.2	6.4	4.3	5.0	3.9	4.8	6.5	5.9	6.2	4.2	5.2	4.3	4.6	6.4
Myeloma	Both Sexes	3.7	3.5	7.1	3.8	1.6	3.1	3.5	3.7	3.5	7.2	3.3	1.7	3.3	3.5
	Male	4.6	4.4	8.5	4.2	1.9	3.7	4.4	4.5	4.4	8.8	3.2	2.0	4.0	4.4
	Female	3.1	2.8	6.3	3.5	1.5	2.6	2.8	3.0	2.9	6.2	3.5	1.5	2.9	2.9
Leukemia	Both Sexes	7.5	7.7	6.7	4.8	3.9	5.1	7.8	7.3	7.6	6.9	3.9	4.1	5.0	7.8
	Male	10.0	10.3	8.8	5.6	5.0	6.4	10.4	9.7	10.2	9.2	4.9	5.3	6.3	10.4
	Female	5.7	5.8	5.3	4.1	3.0	4.1	5.9	5.6	5.8	5.4	2.8	3.2	4.0	5.9

- ^a NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- ^b The SEER 17 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey.
- ^c Rates for American Indian/Alaska Native (AI/AN) are based on the CHSDA(Contract Health Service Delivery Area) counties.
- ^d Asian/Pacific Islander.
- ^e Hispanic (Hisp) and White Non-Hispanic (W-NHisp) are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Mortality data for Hispanics and White Non-Hispanics do not include cases from Minnesota, New Hampshire, and North Dakota.
- Statistic could not be calculated due to less than 16 cases in the time interval.