

Table 17.1

Mesothelioma

Age-Adjusted SEER Cancer Incidence Rates^a, 1980-2007

By Race, Sex and Year of Diagnosis

Race/Sex	Year of Diagnosis													
	1980- 1981	1982- 1983	1984- 1985	1986- 1987	1988- 1989	1990- 1991	1992- 1993	1994- 1995	1996- 1997	1998- 1999	2000- 2001	2002- 2003	2004- 2005	2006- 2007
All Races														
Both Sexes	0.9	0.9	1.0	1.0	1.1	1.1	1.1	1.2	1.1	1.1	1.1	1.0	1.1	1.0
Males														
All Ages	1.6	1.6	1.9	1.7	2.0	2.1	2.3	2.2	2.0	2.1	2.0	1.8	2.0	1.7
Ages 0-54	0.3	0.3	0.3	0.2	0.3	0.2	0.2	0.3	0.2	0.2	0.1	0.1	0.1	0.1
Ages 55-64	3.7	3.7	4.5	3.5	3.1	4.4	3.7	3.3	4.2	3.7	2.6	2.9	2.7	2.4
Ages 65-74	8.2	7.2	8.6	8.3	11.8	9.6	10.6	10.3	9.0	9.8	9.3	7.7	8.5	7.8
Ages 75-84	10.0	11.7	12.0	12.6	12.7	17.2	16.1	17.9	15.2	16.8	19.5	17.3	18.0	14.3
Ages 85+	-	-	-	-	12.8	12.0	20.5	12.4	15.8	15.9	14.1	12.9	18.7	14.3
Females														
All Ages	0.3	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4
Ages 0-54	-	0.1	-	0.1	0.1	0.1	-	0.1	0.1	0.2	0.1	0.1	0.1	0.1
Ages 55-64	0.9	1.0	0.8	1.0	-	0.9	0.9	1.2	0.8	-	-	0.9	0.8	0.5
Ages 65-74	-	2.1	1.4	1.4	1.7	1.4	1.3	1.6	1.5	1.6	1.2	1.4	1.5	1.8
Ages 75-84	-	2.0	3.1	2.3	1.7	3.0	2.8	2.6	2.3	2.0	2.8	2.7	3.2	3.8
Ages 85+	-	-	-	-	-	-	-	-	-	-	2.9	-	-	-
Whites														
Both Sexes	0.9	1.0	1.1	1.0	1.1	1.2	1.2	1.2	1.2	1.2	1.2	1.1	1.2	1.1
Males	1.7	1.7	2.0	1.8	2.2	2.3	2.5	2.3	2.2	2.3	2.3	2.0	2.2	1.9
Females	0.3	0.5	0.4	0.5	0.4	0.5	0.4	0.5	0.4	0.4	0.4	0.4	0.5	0.5
Blacks														
Both Sexes	-	0.6	0.6	0.6	0.7	0.7	0.7	0.9	0.7	0.6	0.4	0.4	0.5	0.3
Males	-	-	-	-	-	1.5	1.3	1.8	1.1	1.3	-	-	0.8	-
Females	-	-	-	-	-	-	-	-	-	-	-	-	-	-

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
 - Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 17.2
MesotheliomaAge-adjusted SEER Incidence^a Rates by Year, Race and Sex

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
Year of Diagnosis:									
1975	0.63	1.00	0.31	0.63	1.00	0.33	-	-	-
1976	0.67	1.13	0.35	0.71	1.21	0.36	-	-	-
1977	0.62	1.05	0.31	0.65	1.06	0.34	-	-	-
1978	0.73	1.37	0.25	0.79	1.50	0.25	-	-	-
1979	0.78	1.51	0.27	0.80	1.60	0.25	-	-	-
1980	0.98	1.83	0.35	1.05	1.97	0.39	-	-	-
1981	0.75	1.42	0.27	0.80	1.50	0.29	-	-	-
1982	0.83	1.42	0.38	0.90	1.53	0.42	-	-	-
1983	1.05	1.86	0.49	1.09	1.95	0.51	-	-	-
1984	1.06	2.04	0.36	1.15	2.24	0.37	-	-	-
1985	0.92	1.67	0.37	0.97	1.73	0.39	-	-	-
1986	0.94	1.67	0.43	1.01	1.79	0.49	-	-	-
1987	0.97	1.73	0.43	1.03	1.82	0.47	-	-	-
1988	0.98	1.85	0.39	1.05	1.98	0.41	-	-	-
1989	1.13	2.20	0.37	1.20	2.35	0.39	-	-	-
1990	1.15	2.15	0.46	1.25	2.32	0.50	-	-	-
1991	1.11	2.10	0.42	1.18	2.21	0.46	-	-	-
1992	1.23	2.48	0.39	1.33	2.71	0.42	-	-	-
1993	1.06	2.04	0.38	1.14	2.24	0.37	-	-	-
1994	1.16	2.17	0.43	1.24	2.32	0.46	1.06	-	-
1995	1.16	2.17	0.46	1.23	2.30	0.48	-	-	-
1996	1.15	2.14	0.45	1.27	2.38	0.50	-	-	-
1997	1.01	1.92	0.37	1.06	2.06	0.35	0.92	-	-
1998	1.08	2.06	0.40	1.18	2.23	0.47	-	-	-
1999	1.12	2.17	0.37	1.20	2.30	0.40	-	-	-
2000	1.08	2.11	0.34	1.22	2.37	0.38	-	-	-
2001	1.06	1.97	0.41	1.17	2.17	0.44	-	-	-
2002	1.02	1.93	0.38	1.15	2.16	0.43	-	-	-
2003	0.94	1.71	0.39	1.04	1.87	0.43	-	-	-
2004	1.10	2.03	0.44	1.21	2.26	0.47	-	-	-
2005	1.03	1.90	0.42	1.18	2.14	0.49	-	-	-
2006	0.95	1.61	0.45	1.05	1.78	0.48	-	-	-
2007	0.96	1.71	0.43	1.08	1.92	0.48	-	-	-
1975-2007	1.00	1.87	0.39	1.08	2.02	0.42	0.56	1.01	0.25

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).
 Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
 - Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 17.3
MesotheliomaSEER Incidence^a Rates, Age-Adjusted and Age-Specific Rates, by Race and Sex

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>SEER Incidence</u>									
<u>Age at Diagnosis</u>									
Age-Adjusted Rates, 2003-2007									
All ages	1.1	1.9	0.4	1.2	2.1	0.5	0.6	1.2	0.3
Under 65	0.3	0.4	0.2	0.3	0.4	0.2	0.2	0.3	0.1
65 and over	6.4	12.5	2.3	7.2	13.7	2.6	3.6	7.5	1.3
All ages (IARC world std) ^b	0.6	1.1	0.3	0.7	1.2	0.3	0.4	0.7	0.2
Age-Specific Rates, 2003-2007									
<1	-	-	-	-	-	-	-	-	-
1-4	-	-	-	-	-	-	-	-	-
5-9	-	-	-	-	-	-	-	-	-
10-14	-	-	-	-	-	-	-	-	-
15-19	-	-	-	-	-	-	-	-	-
20-24	-	-	-	-	-	-	-	-	-
25-29	-	-	-	-	-	-	-	-	-
30-34	-	-	-	-	-	-	-	-	-
35-39	0.1	-	-	0.1	-	-	-	-	-
40-44	0.2	0.2	0.2	0.2	0.2	0.2	-	-	-
45-49	0.3	0.3	0.2	0.3	0.3	0.2	-	-	-
50-54	0.7	0.8	0.5	0.7	0.8	0.6	0.7	-	-
55-59	1.2	1.9	0.6	1.3	2.1	0.6	1.1	1.8	-
60-64	2.4	4.0	1.0	2.7	4.4	1.0	1.6	-	-
65-69	3.6	6.4	1.2	4.1	7.0	1.5	1.8	3.5	-
70-74	5.8	10.3	2.2	6.5	11.4	2.4	3.4	6.4	-
75-79	8.2	15.6	2.8	9.1	17.1	3.3	5.2	11.0	-
80-84	9.7	19.3	3.7	10.8	21.4	4.1	4.7	-	-
85+	7.2	17.0	2.5	7.8	18.5	2.7	4.2	-	-

^a SEER 17 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.^b Rates are per 100,000 and are age-adjusted to the IARC world standard population.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 17.4
Mesothelioma

5-Year Relative and Period Survival (Percent) by Race, Sex, Diagnosis Year, Stage and Age

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
5-Year Relative Survival (Percent)									
Year of Diagnosis:									
1960-1963 ^a	-	-	-	-	-	-	-	-	-
1970-1973 ^a	-	-	-	-	-	-	-	-	-
1975-1977 ^b	10.4	3.2	28.0 ^f	11.2	3.6	28.6 ^f	-	-	-
1978-1980 ^b	6.3	5.8	8.3	6.1	5.4	8.9	-	-	-
1981-1983 ^b	8.2	5.7	15.6	8.6	5.8	16.7	-	-	-
1984-1986 ^b	7.1	4.7	14.7	6.6	4.0	14.9	-	-	-
1987-1989 ^b	6.6	4.8	13.0	6.7	4.9	13.1	4.5	-	-
1990-1992 ^b	8.5	4.6	21.9	8.1	4.3	21.3	12.8 ^f	12.0 ^f	-
1993-1995 ^b	7.1	3.5	19.6	6.4	3.0	18.9	9.6 ^f	8.4 ^f	-
1996-1998 ^b	10.5	8.8	16.3	10.8	8.9	17.4	11.6 ^f	-	-
1999-2006 ^b	7.7	5.6 ^e	14.5	7.1	5.1 ^e	13.9	15.1 ^f	15.4 ^f	-
5-Year Period Survival (Percent) ^{cd}									
Year of Diagnosis:									
1999-2006	7.8	5.9	13.8	7.2	5.3	13.6	11.9	14.2	-
5-Year Relative Survival (Percent), 1999-2006 ^c									
Age at Diagnosis:									
Ages <45	39.7	34.2 ^f	46.0 ^f	36.1	33.3 ^f	39.6 ^f	-	-	-
Ages 45-54	15.7	11.1	22.7	14.8	10.3	21.7	29.7 ^f	-	-
Ages 55-64	7.5	5.5	15.3	7.4	4.8	18.1	3.5	5.4 ^f	-
Ages 65-74	6.0	6.0	6.4	5.9	5.9	6.1	5.1	5.1	-
Ages 75+	2.7	2.0	5.9	2.8	1.9	6.2	0.0	0.0	-
Ages <65	13.3	9.2	23.4	12.4	8.5	22.7	17.7	13.0 ^f	24.7 ^f
Ages 65+	4.2	3.8	6.1	4.2	3.7	6.2	3.4	4.1	0.0

^a Based on End Results data from a series of hospital registries and one population-based registry.^b Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta). Based on follow-up of patients into 2007.^c Based on the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). California excluding SF/SJM/LA, Kentucky, Louisiana, and New Jersey contribute cases for diagnosis years 2000-2006. The remaining 13 SEER Areas contribute cases for the entire period 1999-2006. Based on follow-up of patients into 2007.^d Period survival provides a more up-to-date estimate of survival by piecing together the most recent conditional survival estimates from several cohorts. It is computed here using three year calendar blocks (2004-2006: 0-1 year survival), (2003-2005: 1-2 year survival), (2002-2004: 2-3 year survival), (2001-2003: 3-4 year survival), (2000-2002: 4-5 years survival).^e The difference between 1975-1977 and 1999-2006 is statistically significant ($p < .05$).^f The standard error is between 5 and 10 percentage points.^g The standard error is greater than 10 percentage points.

- Statistic could not be calculated due to fewer than 25 cases during the time period.

Table 17.5
MesotheliomaSEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

	Year of Diagnosis																			
	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
Survival Time																				
1-year	37.4	38.9	37.9	40.8	35.5	40.9	33.3	39.0	37.0	43.1	37.0	39.5	33.9	41.6	43.1	32.8	48.2	39.5	45.3	46.1
2-year	16.7	19.6	17.2	16.1	13.9	20.3	14.1	17.9	14.0	26.4	17.2	22.7	19.0	15.3	25.3	17.2	22.0	16.8	21.3	
3-year	11.2	12.3	11.2	10.8	11.9	13.1	9.1	12.7	12.1	17.9	10.7	15.8	13.1	10.5	15.3	7.2	16.1	10.4		
4-year	9.8	8.9	8.5	6.2	9.0	11.0	6.4	10.1	9.3	12.3	8.3	13.2	11.0	7.0	11.9	5.0	12.8			
5-year	8.4	7.6	7.1	6.2	8.8	9.8	4.4	8.9	7.6	10.6	6.7	13.2	9.2	3.0	11.8	4.6				
6-year	7.2	6.4	5.7	4.3	8.0	7.4	4.4	8.7	6.5	10.6	5.2	12.2	8.4	1.3	10.9					
7-year	6.9	5.9	5.0	3.2	7.7	7.1	4.4	8.2	5.6	9.3	3.4	11.9	8.0	1.3						
8-year	6.4	5.7	4.6	2.7	7.1	7.1	3.7	7.6	5.6	7.2	3.4	9.6	8.0							
9-year	6.2	5.5	4.5	2.7	5.2	7.1	3.7	7.6	4.4	5.5	3.0	9.6								
10-year	6.2	5.3	4.3	2.7	5.2	6.2	3.3	7.6	3.7	5.0	3.0									
11-year	6.2	5.3	4.3	2.7	5.0	5.8	3.3	7.6	3.7	4.6										
12-year	6.1	5.3	4.3	2.7	5.0	5.1	3.3	7.2	3.5											
13-year	6.0	5.3	4.3	2.7	5.0	5.1	2.9	6.7												
14-year	6.0	5.3	4.3	2.7	5.0	5.1	2.9													
15-year	6.0	5.2	4.3	2.7	4.1	5.1														
16-year	5.7	5.2	4.3	2.7	4.1															
17-year	5.7	5.2	4.3	2.7																
18-year	5.7	5.2	4.3																	
19-year	5.7	5.2																		
20-year	5.7	5.2																		

^a

Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 17.6

Mesothelioma

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
Survival Time																				
1-year	35.8	37.5	36.2	37.2	31.4	36.6	35.4	34.6	36.3	46.5	34.3	35.3	32.0	42.0	39.1	31.9	47.4	38.4	45.0	42.3
2-year	12.0	16.5	14.8	12.8	8.6	15.4	14.0	11.5	11.2	26.0	14.6	18.6	14.7	14.3	21.7	16.3	18.6	15.8	17.5	
3-year	6.7	9.8	8.7	7.8	7.9	8.1	8.0	7.0	9.2	16.4	9.0	12.8	7.7	8.9	12.2	6.2	13.3	9.2		
4-year	5.7	7.0	6.2	4.4	4.3	6.4	5.3	4.0	6.2	11.4	6.6	10.6	5.4	6.5	8.9	3.2	12.2			
5-year	4.2	5.5	4.9	4.4	4.3	4.8	2.4	3.5	4.4	10.5	4.6	10.6	4.8	3.8	8.5	3.2				
6-year	3.5	4.7	3.4	3.3	4.3	2.9	2.4	3.5	3.7	10.5	2.4	8.7	4.8	1.6	7.8					
7-year	3.1	4.0	2.8	1.7	4.3	2.3	2.4	3.0	2.3	9.9	1.7	8.7	4.8	1.6						
8-year	2.5	3.8	2.8	0.9	4.3	2.3	1.0	2.4	2.3	7.1	1.7	5.4	4.8							
9-year	2.3	3.8	2.6	0.9	2.9	2.3	1.0	2.4	0.9	5.6	0.9	5.4								
10-year	2.3	3.8	2.4	0.9	2.9	1.9	1.0	2.4	0.0	4.9	0.9									
11-year	2.3	3.8	2.4	0.9	2.2	1.9	1.0	2.4	0.0	4.2										
12-year	2.3	3.8	2.4	0.9	2.2	1.9	1.0	2.1	0.0											
13-year	2.0	3.8	2.4	0.9	2.2	1.9	1.0	2.1												
14-year	2.0	3.8	2.4	0.9	2.2	1.9	1.0													
15-year	1.4	3.8	2.3	0.9	2.2	1.9														
16-year	1.4	3.8	2.3	0.9	2.2															
17-year	1.4	3.8	2.3	0.0																
18-year	1.4	3.8	2.3																	
19-year	1.4	3.8																		
20-year	1.4	3.8																		

^a

Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 17.7
MesotheliomaSEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
Survival Time																				
1-year	42.5	43.9	44.0	53.1	50.7	60.3	25.9	57.2	39.2	32.6	48.6	54.8	41.9	39.7	54.6	36.1	50.6	43.2	46.0	56.5
2-year	31.5	29.9	25.3	27.2	33.1	41.8	14.5	44.3	23.6	27.3	27.8	37.7	37.5	19.2	35.6	20.0	32.0	20.0	33.7	
3-year	24.9	20.5	19.4	20.9	24.6	34.2	12.3	35.3	21.7	22.2	17.4	26.8	35.8	17.1	24.0	10.3	23.7	13.9		
4-year	22.2	14.9	16.0	11.9	24.6	30.3	10.3	33.9	20.1	14.9	14.7	22.6	34.0	8.8	20.3	10.3	14.6			
5-year	20.9	13.9	14.1	11.9	23.6	30.3	10.3	29.6	18.5	11.0	14.7	22.6	26.8	0.0	20.3	8.4				
6-year	18.1	11.3	12.9	7.5	19.0	26.2	10.3	28.0	16.2	9.2	14.7	22.6	21.5	0.0	19.1					
7-year	17.4	11.3	11.9	7.5	16.7	26.2	10.3	28.0	16.2	7.1	10.2	22.0	18.9	0.0						
8-year	17.1	11.3	9.8	7.5	14.4	26.2	10.3	26.5	16.2	7.1	10.2	22.0	18.9							
9-year	17.1	10.4	9.8	7.5	12.6	25.4	10.3	26.5	15.8	5.2	10.2	22.0								
10-year	16.2	8.6	9.8	7.5	12.6	22.8	10.3	26.3	15.8	5.2	10.2									
11-year	15.7	8.6	9.8	7.5	12.6	19.7	10.3	26.3	15.8	5.2										
12-year	15.7	8.6	9.8	7.5	12.6	16.0	10.3	24.9	14.8											
13-year	15.7	8.6	9.8	7.5	12.6	16.0	7.9	22.1												
14-year	15.7	8.6	9.8	7.5	12.6	16.0	7.9													
15-year	15.7	8.6	9.8	7.5	8.8	16.0														
16-year	15.7	8.6	9.8	7.5	8.8															
17-year	15.6	8.6	9.8	7.5																
18-year	15.6	8.6	9.8																	
19-year	15.6	8.6																		
20-year	15.6	8.6																		

^a

Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table 17.8
Mesothelioma (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,
Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and
Lifetime Risk of Dying from Cancer Given Alive at Current Age
Both Sexes, 2005-2007 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer
		+10 yrs	+20 yrs	+30 yrs	Ever				+10 yrs	+20 yrs	+30 yrs	Ever	
All Races	0	0.00	0.00	0.00	0.12	0.10	Asian/ Pacific Islander	0	0.00	0.00	0.00	0.05	0.03
	10	0.00	0.00	0.00	0.13	0.10		10	0.00	0.00	0.00	0.05	0.03
	20	0.00	0.00	0.00	0.13	0.10		20	0.00	0.00	0.00	0.05	0.03
	30	0.00	0.00	0.01	0.13	0.10		30	0.00	0.00	0.01	0.05	0.03
	40	0.00	0.01	0.04	0.13	0.10		40	0.00	0.01	0.01	0.05	0.03
	50	0.01	0.03	0.08	0.13	0.10		50	0.00	0.01	0.03	0.05	0.03
	60	0.03	0.08	0.12	0.13	0.10		60	0.01	0.02	0.04	0.05	0.02
	70	0.06	0.11	-	0.11	0.09		70	0.01	0.04	-	0.04	0.02
White	80	0.07	-	-	0.08	0.06	American Indian/ Alaska Native ^a	80	0.03	-	-	0.03	0.01
	0	0.00	0.00	0.00	0.14	0.11		0	0.00	0.00	0.00	0.07	0.06
	10	0.00	0.00	0.00	0.14	0.11		10	0.00	0.00	0.00	0.07	0.06
	20	0.00	0.00	0.00	0.14	0.11		20	0.00	0.00	0.01	0.07	0.06
	30	0.00	0.00	0.01	0.14	0.11		30	0.00	0.01	0.02	0.07	0.06
	40	0.00	0.01	0.04	0.14	0.11		40	0.01	0.02	0.04	0.07	0.06
	50	0.01	0.04	0.09	0.14	0.11		50	0.02	0.03	0.05	0.07	0.06
	60	0.03	0.09	0.13	0.14	0.11		60	0.02	0.04	0.05	0.05	0.06
Black	70	0.06	0.12	-	0.13	0.10	Hispanic ^b	70	0.02	0.04	-	0.04	0.05
	80	0.07	-	-	0.08	0.07		80	0.03	-	-	0.03	0.02
	0	0.00	0.00	0.00	0.06	0.04		0	0.00	0.00	0.00	0.12	0.06
	10	0.00	0.00	0.00	0.06	0.04		10	0.00	0.00	0.00	0.12	0.06
	20	0.00	0.00	0.00	0.06	0.04		20	0.00	0.00	0.00	0.12	0.06
	30	0.00	0.00	0.01	0.06	0.04		30	0.00	0.00	0.01	0.12	0.06
	40	0.00	0.01	0.02	0.06	0.04		40	0.00	0.01	0.03	0.12	0.06
	50	0.01	0.02	0.05	0.07	0.04		50	0.01	0.03	0.07	0.12	0.06
	60	0.02	0.04	0.06	0.06	0.04		60	0.02	0.06	0.11	0.12	0.06
	70	0.03	0.05	-	0.06	0.04		70	0.05	0.10	-	0.11	0.06
	80	0.03	-	-	0.03	0.03		80	0.07	-	-	0.08	0.04

Devcan Version 6.5.0, June 2010, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

Underlying mortality data for Hispanics exclude deaths from the District of Columbia, Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.

Table 17.9
Mesothelioma (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,
Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and
Lifetime Risk of Dying from Cancer Given Alive at Current Age
Males, 2005-2007 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer
		+10 yrs	+20 yrs	+30 yrs	Ever				+10 yrs	+20 yrs	+30 yrs	Ever	
All Races	0	0.00	0.00	0.00	0.20	0.16	Asian/ Pacific Islander	0	0.00	0.00	0.00	0.08	0.04
	10	0.00	0.00	0.00	0.21	0.17		10	0.00	0.00	0.00	0.09	0.04
	20	0.00	0.00	0.00	0.21	0.17		20	0.00	0.00	0.00	0.09	0.04
	30	0.00	0.00	0.02	0.21	0.17		30	0.00	0.00	0.01	0.09	0.04
	40	0.00	0.02	0.06	0.21	0.17		40	0.00	0.01	0.02	0.09	0.04
	50	0.01	0.05	0.13	0.22	0.18		50	0.00	0.02	0.05	0.09	0.04
	60	0.04	0.13	0.20	0.22	0.18		60	0.02	0.04	0.08	0.09	0.04
	70	0.10	0.19	-	0.21	0.17		70	0.03	0.07	-	0.08	0.04
White	80	0.13	-	-	0.15	0.13	American Indian/ Alaska Native ^a	80	0.05	-	-	0.06	0.03
	0	0.00	0.00	0.00	0.22	0.18		0	0.00	0.00	0.00	0.10	0.07
	10	0.00	0.00	0.00	0.23	0.18		10	0.00	0.00	0.00	0.10	0.07
	20	0.00	0.00	0.00	0.23	0.18		20	0.00	0.00	0.01	0.10	0.07
	30	0.00	0.00	0.02	0.23	0.19		30	0.00	0.01	0.03	0.11	0.07
	40	0.00	0.02	0.06	0.23	0.19		40	0.01	0.03	0.05	0.11	0.08
	50	0.01	0.06	0.15	0.24	0.19		50	0.02	0.04	0.07	0.11	0.08
	60	0.05	0.15	0.23	0.24	0.20		60	0.02	0.05	0.09	0.09	0.08
Black	70	0.11	0.21	-	0.23	0.19	Hispanic ^b	70	0.04	0.09	-	0.09	0.06
	80	0.14	-	-	0.17	0.14		80	0.07	-	-	0.07	0.01
	0	0.00	0.00	0.00	0.09	0.06		0	0.00	0.00	0.00	0.18	0.10
	10	0.00	0.00	0.00	0.10	0.06		10	0.00	0.00	0.00	0.18	0.10
	20	0.00	0.00	0.00	0.10	0.07		20	0.00	0.00	0.00	0.18	0.10
	30	0.00	0.00	0.01	0.10	0.07		30	0.00	0.00	0.02	0.18	0.11
	40	0.00	0.01	0.04	0.10	0.07		40	0.00	0.01	0.04	0.18	0.11
	50	0.01	0.03	0.08	0.11	0.07		50	0.01	0.04	0.10	0.19	0.11
	60	0.03	0.08	0.10	0.11	0.08		60	0.03	0.10	0.18	0.19	0.11
	70	0.07	0.10	-	0.11	0.07		70	0.08	0.17	-	0.18	0.10
	80	0.06	-	-	0.07	0.06		80	0.13	-	-	0.14	0.07

Devcan Version 6.5.0, June 2010, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

Underlying mortality data for Hispanics exclude deaths from the District of Columbia, Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.

Table 17.10
Mesothelioma (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,
Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and
Lifetime Risk of Dying from Cancer Given Alive at Current Age
Females, 2005-2007 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer
		+10 yrs	+20 yrs	+30 yrs	Ever				+10 yrs	+20 yrs	+30 yrs	Ever	
All Races	0	0.00	0.00	0.00	0.06	0.04	Asian/ Pacific Islander	0	0.00	0.00	0.00	0.02	0.01
	10	0.00	0.00	0.00	0.06	0.04		10	0.00	0.00	0.00	0.02	0.01
	20	0.00	0.00	0.00	0.06	0.04		20	0.00	0.00	0.00	0.02	0.01
	30	0.00	0.00	0.01	0.06	0.04		30	0.00	0.00	0.00	0.02	0.01
	40	0.00	0.01	0.02	0.06	0.04		40	0.00	0.00	0.01	0.02	0.01
	50	0.01	0.02	0.04	0.06	0.04		50	0.00	0.01	0.01	0.02	0.01
	60	0.01	0.03	0.05	0.05	0.03		60	0.00	0.01	0.01	0.01	0.01
	70	0.02	0.04	-	0.05	0.03		70	0.00	0.01	-	0.01	0.01
White	80	0.03	-	-	0.03	0.02	American Indian/ Alaska Native ^a	80	0.01	-	-	0.01	0.00
	0	0.00	0.00	0.00	0.06	0.04		0	0.00	0.00	0.00	0.04	0.04
	10	0.00	0.00	0.00	0.06	0.04		10	0.00	0.00	0.00	0.04	0.04
	20	0.00	0.00	0.00	0.06	0.04		20	0.00	0.00	0.01	0.04	0.04
	30	0.00	0.00	0.01	0.06	0.04		30	0.00	0.01	0.02	0.04	0.04
	40	0.00	0.01	0.02	0.06	0.04		40	0.01	0.02	0.03	0.04	0.04
	50	0.01	0.02	0.04	0.06	0.04		50	0.01	0.03	0.03	0.03	0.04
	60	0.01	0.04	0.05	0.06	0.04		60	0.02	0.02	0.02	0.02	0.04
Black	70	0.03	0.05	-	0.05	0.03	Hispanic ^b	70	0.00	0.00	-	0.00	0.03
	80	0.03	-	-	0.03	0.02		80	0.00	-	-	0.00	0.03
	0	0.00	0.00	0.00	0.03	0.01		0	0.00	0.00	0.00	0.07	0.03
	10	0.00	0.00	0.00	0.03	0.01		10	0.00	0.00	0.00	0.07	0.03
	20	0.00	0.00	0.00	0.03	0.01		20	0.00	0.00	0.00	0.07	0.03
	30	0.00	0.00	0.01	0.03	0.01		30	0.00	0.00	0.01	0.07	0.03
	40	0.00	0.01	0.02	0.03	0.01		40	0.00	0.01	0.02	0.07	0.03
	50	0.01	0.02	0.02	0.03	0.01		50	0.01	0.02	0.04	0.07	0.03
	60	0.01	0.02	0.03	0.03	0.01		60	0.01	0.03	0.06	0.07	0.02
	70	0.01	0.02	-	0.02	0.01		70	0.02	0.05	-	0.06	0.02
	80	0.01	-	-	0.02	0.01		80	0.03	-	-	0.04	0.02

Devcan Version 6.5.0, June 2010, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

Underlying mortality data for Hispanics exclude deaths from the District of Columbia, Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.

Table 17.11
Mesothelioma

SEER Incidence
Age-Adjusted Rates and Trends^a
By Race/Ethnicity and Sex

SEER Incidence RACE/ETHNICITY	SEER 17 Areas ^b Rate 2003-2007 Rate per 100,000 persons			SEER 13 Areas ^b Trend 1998-2007 ^c AAPC (%)			SEER 9 Areas ^b Trend 1998-2007 ^d AAPC (%)		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
All Races	1.1	1.9	0.4	-1.1*	-1.6*	1.0	-1.2*	-1.8*	0.0
White	1.2	2.1	0.5	-0.7*	-1.3*	0.8*	-0.7	-1.5*	0.9*
White Hispanic ^e	1.0	1.6	0.5	-2.5*	-3.2*	-	-	-	-
White Non-Hispanic ^e	1.2	2.2	0.5	-0.6*	-1.1*	0.6	-	-	-
Black	0.6	1.2	0.3	-	-	-	-7.0*	-7.9*	-
Asian/Pacific Islander	0.4	0.6	0.2	-	-	-	-	-	-
Amer Ind/Alaska Nat ^f	0.8	-	-	-	-	-	-	-	-
Hispanic ^e	0.9	1.5	0.5	-2.3*	-3.0*	-	-	-	-

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 3.4.3, April 2010, National Cancer Institute.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry and Rural Georgia.

The SEER 17 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey.

^c The 1998-2007 AAPC estimates are based on a Joinpoint analysis with up to 2 Joinpoints over diagnosis years 1992-2007.

^d The 1998-2007 AAPC estimates are based on a Joinpoint analysis with up to 4 Joinpoints over diagnosis years 1975-2007.

^e Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

^f Incidence data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

* The APC is significantly different from zero (p<.05).

Table 17.12
MesotheliomaAge-Adjusted SEER Incidence Rates^a
By Registry, Race and Sex

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>SEER Incidence Rates^a, 2003-2007</u>									
Atlanta & Rural Georgia	0.55	1.05	0.23	0.67	1.22	-	-	-	-
Atlanta	0.55	1.06	0.24	0.69	1.24	-	-	-	-
Rural Georgia	-	-	-	-	-	-	-	-	-
California	0.97	1.77	0.39	1.10	1.99	0.44	0.62	1.18	-
Greater Bay Area	0.97	1.82	0.34	1.20	2.25	0.41	-	-	-
San Francisco-Oakland	0.99	1.86	0.35	1.24	2.31	0.43	-	-	-
San Jose-Monterey	0.93	1.74	0.32	1.14	2.16	0.38	-	-	-
Los Angeles	0.88	1.57	0.40	1.05	1.85	0.49	0.58	1.15	-
Greater California	1.02	1.85	0.40	1.10	1.98	0.43	0.78	1.53	-
Connecticut	1.14	2.02	0.53	1.18	2.12	0.52	-	-	-
Detroit	0.98	1.75	0.47	1.12	2.04	0.50	0.41	-	-
Hawaii	0.67	1.26	-	1.04	-	-	-	-	-
Iowa	0.77	1.37	0.34	0.77	1.40	0.34	-	-	-
Kentucky	0.75	1.38	0.32	0.77	1.39	0.33	-	-	-
Louisiana	1.35	2.50	0.53	1.45	2.64	0.58	1.12	2.17	-
New Jersey	1.52	2.89	0.61	1.65	3.12	0.65	0.78	1.45	-
New Mexico	0.97	1.49	0.54	0.96	1.45	0.57	-	-	-
Seattle-Puget Sound	1.52	2.75	0.61	1.62	2.95	0.61	-	-	-
Utah	0.90	1.52	0.38	0.93	1.57	0.39	-	-	-
9 SEER Areas ^b	1.00	1.79	0.43	1.11	1.99	0.47	0.41	0.70	0.22
11 SEER Areas ^b	0.97	1.74	0.41	1.10	1.97	0.47	0.45	0.81	0.23
13 SEER Areas ^b	0.96	1.74	0.41	1.10	1.97	0.47	0.46	0.82	0.22
17 SEER Areas ^b	1.06	1.93	0.44	1.17	2.11	0.48	0.65	1.21	0.29

^a

Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130)

^b

The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta.

The SEER 11 areas comprise the SEER 9 areas plus San Jose-Monterey and Los Angeles.

The SEER 13 areas comprise the SEER 11 areas plus the Alaska Native Registry and Rural Georgia.

The SEER 17 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey.

-

Statistic not shown. Rate based on less than 16 cases for the time interval.

Table 17.13
MesotheliomaEstimated United States Cancer Prevalence Counts^a on January 1, 2007
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <17 ^e	0 to <32 ^e	>=32 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	2,943	596	320	238	150	159	3,965	4,427	+	+
	Males	2,066	286	105	58	80	60	2,489	2,655	+	+
	Females	877	310	215	180	70	99	1,476	1,772	+	+
White ^b	Both Sexes	2,749	485	283	235	127	136	3,624	4,037	+	+
	Males	1,934	231	92	55	57	47	2,290	2,417	+	+
	Females	815	254	191	180	70	89	1,334	1,620	+	+
Black ^b	Both Sexes	97	69	26	0	22	24	191	236	+	+
	Males	61	34	13	0	22	13	107	142	+	+
	Females	36	35	13	0	0	11	84	94	+	+
Asian/ Pacific Islander ^c	Both Sexes	47	12	5	+	+	+	64	+	+	+
	Males	35	9	5	+	+	+	49	+	+	+
	Females	12	3	0	+	+	+	15	+	+	+
Hispanic ^d	Both Sexes	261	62	10	+	+	+	338	+	+	+
	Males	153	46	0	+	+	+	199	+	+	+
	Females	108	16	10	+	+	+	139	+	+	+

Estimated prevalence percent^a on January 1, 2007, of the SEER 11 population diagnosed in the previous 17 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)										Age-Adjusted ^f
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
Race	Sex											
All Races ^c	Both Sexes	0.0012%	-	-	-	0.0002%	0.0006%	0.0016%	0.0038%	0.0063%	0.0066%	0.0012%
	Males	0.0015%	-	-	-	-	0.0004%	0.0019%	0.0057%	0.0106%	0.0124%	0.0018%
	Females	0.0008%	-	-	-	0.0003%	0.0008%	0.0013%	0.0020%	0.0030%	0.0034%	0.0008%
White ^c	Both Sexes	0.0013%	-	-	-	0.0002%	0.0006%	0.0019%	0.0041%	0.0073%	0.0070%	0.0013%
	Males	0.0017%	-	-	-	-	0.0004%	0.0021%	0.0060%	0.0119%	0.0127%	0.0019%
	Females	0.0010%	-	-	-	0.0003%	0.0008%	0.0016%	0.0023%	0.0036%	0.0039%	0.0009%
Black ^c	Both Sexes	0.0005%	-	-	-	-	-	-	0.0028%	0.0032%	0.0054%	0.0007%
	Males	0.0008%	-	-	-	-	-	-	0.0041%	-	-	0.0012%
	Females	0.0003%	-	-	-	-	-	-	-	-	-	0.0004%
Asian/ Pacific Islander ^c	Both Sexes	0.0005%	-	-	-	-	-	-	0.0016%	-	0.0041%	0.0005%
	Males	0.0008%	-	-	-	-	-	-	0.0036%	-	0.0087%	0.0010%
	Females	0.0002%	-	-	-	-	-	-	-	-	-	0.0002%
Hispanic ^d	Both Sexes	0.0008%	-	-	-	-	0.0007%	0.0020%	0.0036%	0.0078%	0.0095%	0.0014%
	Males	0.0009%	-	-	-	-	-	0.0027%	0.0067%	0.0092%	0.0117%	0.0018%
	Females	0.0006%	-	-	-	-	-	0.0014%	-	0.0068%	0.0082%	0.0011%


- ^a US 2007 cancer prevalence counts are based on 2007 cancer prevalence proportions from the SEER registries and 1/1/2007 US population estimates based on the average of 2006 and 2007 population estimates from the US Bureau of the Census.
- ^{b c d} Prevalence was calculated using the First Malignant Primary Only for a person. Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.
- ^e Maximum limited-duration prevalence: 32 years for 1975-2007 SEER 9 data; 17 years for 1990-2007 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).
- ^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.
- ^{g h i} (g) Cases diagnosed more than 32 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <32 and >=32. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.
- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.
- + Not available.

Figure 17.1

Mesothelioma

SEER Incidence, 1975-2007

All Races, By Sex



Source: SEER 9 areas. Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103). Regression lines are calculated using the Joinpoint Regression Program Version 3.4.3, April 2010, National Cancer Institute.